

Soleto®

MAPP 20225

GROUP 5 HERBICIDE

Pre-emergence herbicide for the control of annual broad-leaved weeds and annual grasses in potatoes.

Suspension concentrate (SC) formulation containing 500 g/L metobromuron.

HERBICIDE

H

Net content

5L

PROTECT FROM FROST

Production date /
Batch number:
see packaging



SOLETO

A suspension concentrate (SC) formulation containing 500 g/L metobromuron.

WARNING

H351 Suspected of causing cancer.

H373 May cause damage to blood system through prolonged or repeated exposure.

H410 Very toxic to aquatic life with long lasting effects.

P273 Avoid release to the environment.

P280 Wear protective gloves/protective clothing.

P301+P312 IF SWALLOWED: Call a POISON CENTRE, doctor if you feel unwell.

P308+P313 IF exposed or concerned: Get medical advice/attention.

P391 Collect spillage.

UFI: KQ46-80HS-5001-055F

P501 Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for empty, clean containers which can be disposed of as non-hazardous waste.

EUH208 Contains metobromuron and 1,2-benzisothiazolin-3-one. May produce an allergic reaction.

EUH401 To avoid risks to human health and the environment, comply with the instructions for use.

For advice on medical emergencies, fires, spillages or chemical hazards ONLY – phone Carechem 24 EU: 01235 239670

For technical and non-emergency calls phone 0845 373 0305

Authorisation Holder:
Certis Belchim BV
Stadsplateau 16
3521 AZ Utrecht
PO Box 607
3500 AP Utrecht
Netherlands

® Registered Trademark of Certis Belchim BV

05012601_2523 • 106044188:UK



Marketing Company: **Certis Belchim BV**
Suite 5, 3 Riverside, Granta Park, Great Abington
Cambridgeshire CB21 6AD
Tel.: 01223 652500 | info.uk@certisbelchim.com
www.certisbelchim.co.uk

IMPORTANT INFORMATION

FOR PROFESSIONAL USE ONLY AS AN AGRICULTURAL HERBICIDE

Crop	Maximum individual dose (L product/ha)	Maximum number of treatments (per crop)	Latest time of application
Potato	4	1	Pre-emergence (BBCH 00-09)

Other specific restriction:

Non returnable containers must not be re-used for any purpose.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER WHICH IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

THE CONTROL OF SUBSTANCES HAZARDOUS TO HEALTH (COSHH) REGULATIONS MAY APPLY TO THE USE OF THIS PRODUCT AT WORK.

SAFETY PRECAUTIONS

Operator Protection

ENGINEERING CONTROL OF OPERATOR EXPOSURE must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE GLOVES when handling the concentrate or handling contaminated surfaces.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide equal or higher standard of protection.

IN CASE OF CONTACT WITH EYES OR SKIN, rinse immediately with plenty of water and consult a specialist.

IN CASE OF ACCIDENT OR IF YOU FEEL UNWELL; seek medical advice immediately (show the label where possible).

DO NOT BREATHE spray vapours or mist.

WASH ALL PROTECTIVE CLOTHING thoroughly after use, especially the inside of gloves.

WASH HANDS AND EXPOSED skin before eating, drinking or smoking and after work.

WHEN USING, DO NOT EAT, DRINK OR SMOKE

Environmental Protection:

DO NOT CONTAMINATE WATER with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

Take care to avoid drift onto neighbouring crops and non-agricultural land.

Storage and Disposal

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

Store in a safe, dry, frost-free place designated as an agrochemical store.

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDINGSTUFFS.

Non-returnable containers must not be re-used for any purpose.



DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

SOLETO® is a selective pre-emergence herbicide for the control of annual broad-leaved weeds and annual grasses in potatoes. The product is mainly absorbed by the roots and transported acropetally within the plant after uptake.

SOLETO acts by inhibiting photosynthesis in susceptible weeds.

SOLETO may be applied on all varieties of potatoes.

WEEDS CONTROLLED

The following weeds are susceptible to SOLETO when applied pre-emergence of the weeds:

Common name of weed	Latin name	Susceptibility at 3 L/ha	Susceptibility at 4 L/ha
Grass weeds			
Annual meadow grass	<i>Poa annua</i>	S	S
Cockspur grass	<i>Echinochloa crus-galli</i>	-	MS
Foxtail spp	<i>Setaria</i> spp	-	MS
Broad-leaved weeds			
Annual mercury	<i>Mercurialis annua</i>	-	MS
Black bindweed	<i>Fallopia convolvulus</i>	MR	MS
Black nightshade	<i>Solanum nigrum</i>	-	MS
Charlock	<i>Sinapis arvensis</i>	S	S
Chickweed	<i>Stellaria media</i>	S	S
Common amaranth	<i>Amaranthus retroflexus</i>	-	MS
Fat hen	<i>Chenopodium album</i>	S	S
Field pansy	<i>Viola arvensis</i>	-	S
Fumitory	<i>Fumaria officinalis</i>	-	MS
Green amaranth	<i>Amaranthus hybridus</i>	-	MS
Groundsel, Common	<i>Senecio vulgaris</i>	S	S
Knotgrass	<i>Polygonum aviculare</i>	MS	S
Scented mayweed	<i>Matricaria chamomilla</i>	S	S
Scentless mayweed	<i>Matricaria inodora</i>	S	S
Shepherd's purse	<i>Capsella bursa-pastoris</i>	S	S

RESTRICTIONS

Some transient crop discolouration may occur under certain climatic conditions where heavy rainfall follows application. This transient crop effect is rapidly outgrown and has no effect on crop yield.

Reduced levels of weed control may occur if application is made to dry cloddy seedbeds or if a prolonged dry period follows application.

Do not overlap spray swaths.

TIMING OF APPLICATION

SOLETO should be applied following the last ridging and before emergence of the crop and weeds.

DOSE RATE

Apply SOLETO at a dose rate of 3 to 4 litres product/ha (see weed susceptibility table for selection of correct dose).

The higher dose rate should be used on heavy soils with a high clay content.

Maximum number of treatments per crop: One

WATER VOLUME AND SPRAY QUALITY

Apply in 200 to 600 L of water per ha using a MEDIUM spray quality as defined by BCPC.

SOIL TYPE

SOLETO is recommended for use on all soil types. Optimum efficacy is achieved when applied to moist soil. The efficacy of SOLETO may be reduced when applied on soils with a high organic matter content (>10%).

MIXING AND APPLICATION

Before using SOLETO, ensure the spray equipment is clean. Shake the container well before use.

1. Half-fill the spray tank with clean water and commence agitation.
2. Add the required amount of product to the spray tank and complete filling, maintaining agitation until spraying is completed.
3. On emptying the product container, rinse container thoroughly by using an integrated pressure rinsing device or manually rinsing three times when empty. Add washings to sprayer at time of filling and dispose of container safely.
4. On preparation of spray solution, use as soon as practically possible.

Never prepare more spray solution than required.

COMPATIBILITY

Consult Certis Belchim before tank mixing with any adjuvants or other pesticides.

SPRAY TANK CLEAN OUT

All application equipment and contaminated clothing should be thoroughly washed/ rinsed with clean water three times. After each step of washing drain the sprayer, spray out completely. Ensure that all liquid is removed from the sprayer tank, pump and hoses. Remove nozzles, open tank and drain pump to allow free access of air to all parts of the system. Care should be taken not to rinse contaminated washings from application equipment into waste water channels. Contaminated cleaning liquids should be disposed of safely according to local regulations.

Operators should read the sprayer manufacturer's instructions before beginning to wash out sprayers.

RESISTANCE

Metobromuron belongs to the chemical family of ureas and is classified as an inhibitor of photosynthesis.

Spraying the same field repeatedly, with products containing active substances from the same chemical family, or having the same mode of action, can lead to the development of resistance.

To reduce the risk, it is recommended to alternate or to tank mix products containing active substances with complementary activity but from different chemical groups or having different modes of action, not only within one growing season but also throughout the crop rotation.

Strains of some annual grasses (e.g. black-grass, wild-oats, and Italian rye-grass) have developed resistance to herbicides which may lead to poor control. A strategy for preventing and managing such resistance should be adopted. Guidelines have been produced by the Weed Resistance Action Group and copies are available from the HGCA, CPA, your distributor, crop adviser or product manufacturer.

FOLLOWING CROPS

Following normal harvest of a treated crop, Brassicas (oilseed rape, turnip, brassica vegetables) and sugar beet may be sown, providing the soil has been ploughed. No restrictions apply when other crops are sown.

REPLACEMENT CROPS

In the event of crop failure for any reason, maize, beans, peas and carrots may be sown after ploughing. It is not recommended to sow Brassicas (oilseed rape, turnip, brassica vegetables) or sugar beet as a replacement crop after crop failure. Potatoes can be re-planted after minimal cultivation.

CONDITIONS OF SUPPLY

All goods supplied by us are of high grade and we believe them to be suitable, but as we cannot exercise control over their storage, handling, mixing or use, or the weather conditions before, during and after application which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality of fitness for any purpose of our goods are excluded, and no responsibility will be accepted by us or re-sellers for any failure in performance, damage or injury whatsoever arising from their storage, handling application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.

To access the Safety Data Sheet for this product scan the QR code or use the weblink below:



WEBLINK to specific data sheet on manufacturer's website:

www.certisbelchim.co.uk/pdf/MSDS/Soletto_MSDS.pdf

Alternatively, contact your supplier.